5th International Conference on X-Ray Lasers Lund, Sweden, June 10-14, 1996

Target Considerations for Inner-Shell Photo-Ionized X-Ray Lasing

Stephen J. Moon and David C. Eder

Lawrence Livermore National Laboratory, Livermore, CA 94550 USA tel: (510) 424-4856, fax: (510) 423-9208, email: sjmoon@llnl.gov

Abstract

We present target design for the inner-shell photo-ionization (ISPI) x-ray laser scheme and discuss the requirements for the ionizing x-ray source in rise time and flux. An investigation of the rapid rise time of x-ray emission from targets heated by an ultrashort-pulse (USP) high-intensity optical laser was conducted for use as the x-ray source for ISPI x-ray lasing. Results of x-ray rise times from instantaneously heated Au rod targets[1] show little benefit for using optical pulse widths less than 30 fs [2]. Modeling using the hydrodynamic/atomic kinetics code LASNEX of a 30 fs USP driving laser with energy of order 1 J incident on a structured Au target composed of vertical rods with diameter of 50 nm predicts sufficient x rays to produce a gain-length product ≈ 10 in C at 45Å. Collisional ionization to the lower lasing level limits the duration of lasing giving a x-ray laser pulse duration of order 60 fs FWHM[3]. Our previous calculations assumed a specified absorbed energy in a skin depth in the rods. Here we are also presenting results for planar targets where the absorption of the USP laser light is determined using a 1-D wave solver in LASNEX. We consider various angles of incident including the effect of super-thermal electrons. These results are compared to structured targets.

*Work performed under the auspices of the U. S. Department of Energy by Lawrence Livermore National Laboratory under Contract W-7405-ENG-48.

- 1. F. Budnik, G. Kulcsár, L. Zhao, R. Marjoribanks, P. Herman, D. Al-Mawlawi, M. Moskovits, 24th Anomalous Absorption Conference, Pacific Grove, CA, June 1994
- 2. S. J. Moon and D. C. Eder, SPIE Proceedings 2520, San Diego, July 1995.
- S. J. Moon, D. C. Eder, and G. L. Strobel, Proceedings of the 4th International Colloquium on X-Ray Lasers, Williamsberg, VA, May 1994, edited by D. C. Eder and D. L. Matthews (AIP, New York).